

Calcium

FOR WOMEN'S HEALTH



ARTICLE BY
GABBY CAMPBELL, BHSC(NAT)

Known for its important role in skeletal health and the prevention of osteoporosis, a reduced dietary calcium intake and an increased physiological need can have a profound effect on the individual beyond bone health. In this article, we review the functions of calcium within the body and look at dietary and supplemental sources of calcium. We take a closer look at the rationale for supplementation across the lifespan and showcase the role of calcium D-glucarate supplementation, looking at its unique role in hormone regulation.

Calcium is an essential macronutrient found in abundance in the body, with 99% stored in teeth and bone.¹ This mineral is involved in several biochemical processes, including:

- Hormonal secretion
- Nerve impulse transmission
- Muscular function
- Vascular contraction
- Vasodilation
- Intracellular signalling.^{1,2}

Lifespan calcium requirements

Calcium plays an important role in women's health, with intake requirements differing across the lifespan.

Higher doses of calcium are required during pregnancy and breastfeeding³ to:

- Minimise bone mineral loss
- Aid skeletal growth of the child⁴
- Support milk production⁴
- Prevent pre-eclampsia.³

Higher doses are also needed for menopausal and post-menopausal women, as the drop in oestrogen necessary for bone formation causes more bone resorption, increasing the risk of osteoporosis and osteoporotic fractures.⁵

Calcium deficiency

With the wide-ranging role of calcium throughout the body, Figure 1 depicts some of the physical expressions of calcium deficiency.



KEY HIGHLIGHTS

- Calcium is crucial for skeletal health, hormonal secretion, nerve impulse transmission, muscular function, vascular contraction, vasodilation, and intracellular signaling.
- Women's calcium needs vary, with higher doses required during pregnancy, breastfeeding, menopause, and post-menopause.
- Many women do not meet their daily calcium needs through diet alone, necessitating supplementation.
- Common supplemental forms of calcium include calcium carbonate, calcium citrate, and calcium hydroxyapatite, each with specific absorption and tolerance characteristics.
- Calcium D-Glucarate aids in liver detoxification and hormonal regulation, particularly in inhibiting beta-glucuronidase.




Figure 1. Signs of calcium deficiency

Calcium in the diet

Meeting the requirements for calcium can be challenging - 1 in 10 Australians do not meet their daily calcium requirements through diet.⁶ While dietary intake of calcium in childhood is generally sufficient, intake drops at adolescence in Australia.⁷ A staggering 90 per cent of women between the ages of 12-18 years and over 50 years were found to have insufficient dietary calcium in Australia, despite an increased need.⁶ The number of Australians choosing to avoid animal products and switching to plant-based alternatives is increasing and while plant-based milk and dairy alternatives provide a dairy replacement from a taste perspective, unless they are fortified with calcium, they often do not provide the calcium benefits of dairy products.⁷

Generally, dietary intake of calcium is sufficient for most people, provided their diet contains no restrictions. There are, however, some individuals who may benefit from supplementation including individuals who:

- Take corticosteroids
- Diagnosed with osteopenia or osteoporosis
- Follow a vegan or dairy-free diet or those with a lactose intolerance
- Women with amenorrhoea
- Residents of aged / long-term care facilities
- Individuals with gastrointestinal (GI) diseases, given dietary calcium is largely absorbed in the small intestine.^{2,8}

Table 1. Analysis of supplemental forms of calcium

FORM OF CALCIUM	CONSIDERATIONS
Calcium carbonate	Generally well absorbed and tolerated when taken with food, though best avoided by those taking H2 blockers or proton pump inhibitors (PPIs), or individuals with hypochlorhydria as it requires an acidic environment for optimal absorption.
Calcium citrate	Best absorbed and most bioavailable form of calcium when taken with food. It can also be taken without food if needed with little, to no GI disturbance. The best form for patients on PPIs or H2 blockers, those with hypochlorhydria, inflammatory bowel disease or other GI absorption disorders. ⁹
Calcium phosphate / Calcium hydroxyapatite	Provides a higher amount of elemental calcium per tablet, so is best for cost and compliance considerations, ⁹ however, is also the form most associated with GI disturbance. ²
Calcium lactate and Calcium gluconate	Best absorbed and most bioavailable form of calcium when taken with food. It can also be taken without food if needed with little, to no GI disturbance.

Supplemental calcium

The most common forms of supplemental calcium include calcium carbonate and calcium citrate, followed by calcium hydroxyapatite. Several factors for consideration when prescribing are outlined in the table below (See Tables 1 and 2). Regardless of the form chosen, absorption is best when taken with food, in divided doses of 500 mg or less at a time and when given concurrently with vitamin D.⁹

Table 2. Calcium supplementation recommended dosage

CONCERN	DOSAGE
General dietary or total calcium needs for women	1000 mg/day¹²
GI diseases such as coeliac disease	1200 - 1500 mg/day plus 400 IU vitamin D/ day¹³
Menopausal and post-menopausal women over 50 years, for the reduction of osteoporosis and osteoporotic fractures	1000 - 1200 mg/day plus 400 IU - 800 IU vitamin D/day^{2,14}
Pregnancy and lactation	1500 - 2000 mg/day³
Women with amenorrhoea	1200 - 1500 mg/day plus 400 - 1000 IU vitamin D/ day¹⁵
If taking certain medications (e.g. Corticosteroids)	1200 mg/day plus 800 - 2000 IU vitamin D/day¹⁶



CALCIUM D-GLUCARATE

Liver and Hormone Regulation

Calcium D-glucarate is the calcium salt and supplemental form of D-glucaric acid, an endogenously produced compound. Calcium's function is to stabilise glucaric acid, not act as a calcium supplement – glucaric acid is the active ingredient in the supplement.

When ingested, calcium D-glucarate liberates D-glucaric acid which is further metabolised into D-glucaro-1,4-lactone or D-glucaro-6,3-lactone known for their beneficial role in liver support, lipid and hormonal regulation, including oestrogen metabolism.¹⁷ Glucaric acids' primary action is through the inhibition of beta-glucuronidase – a specialised enzyme produced by colonic gut microbiota involved in the interference of phase II liver detoxification - glucuronidation. (See Figure 2)¹⁷

Glucuronidation

Glucuronidation is the detoxification process performed by the liver, responsible for the detoxification and elimination of hormones and exogenous toxins. (See Figure 2) The process of glucuronidation involves the binding of glucuronic acid to form glucuronide conjugates with toxins, in preparation for elimination. Excess beta-glucuronidase has the capacity to deconjugate these compounds, allowing for the reabsorption of toxins. Furthermore, excess beta-glucuronidase is associated with an increased risk for hormone-dependent cancers such as breast, colon, and prostate cancers.¹⁷

Supplemental calcium D-glucarate may be considered as a treatment strategy in addition to lifestyle and dietary modifications.¹⁷ Current research suggest supplemental doses of calcium D-glucaric acid of 1500 - 3000 mg/day, with supplementation not recommended during pregnancy due to the role of calcium D-glucarate in hormone detoxification.¹⁷




Figure 2. The function of calcium D-glucarate in detoxification. Calcium D-glucarate enhances phase 2 glucuronidation by inhibiting beta-glucuronidase

CALCIUM SUPPLEMENTATION FOR SPECIFIC CONDITIONS

Calcium supplementation and cardiovascular disease risk in women

Research on calcium supplementation and cardiovascular disease (CVD) has been divisive, with studies producing contradictory results. In some studies, lower doses of calcium supplementation (600 mg/day) decreased CVD risks,² while others demonstrate a considerable increase with doses of 700 –1000 mg/day.¹⁸ The assumption being that increased calcium may contribute to raised serum calcium levels potentially hastening coronary artery calcification.² With no conclusive outcome, and generally favourable results on CVD in the literature, calcium supplementation is still believed to be beneficial overall.²

Calcium supplementation, osteoporosis, and falls risk in women

Calcium requirements increase for women during and post menopause, as reduced levels of oestrogen precipitate a decrease in the absorption and preservation of calcium. This reduction of calcium absorption and preservation can affect bone mineral density and lead to osteoporosis. When combined with weakened muscles, curvature of the spine or poor postural control experienced with ageing, these women tend to also have a higher risk of falls and fracture.¹⁹

Research supports supplementing with both calcium and vitamin D, as a deficiency can further inhibit calcium absorption in older women to positively impact calcium homeostasis, mitigating these concerns.^{7,20}

Calcium supplement for pre-eclampsia prevention during pregnancy

Pre-eclampsia is a complication that can occur in pregnancy and commonly affects 2-5% of pregnant women. It is a multisystem condition characterised by hypertension (≥ 140 mmHg/ ≥ 90 mmHg) and proteinuria (≥ 300 mg/day) typically presenting after 20 weeks' gestation. Other symptoms may include neurological and haematological complications, in addition to liver abnormalities and acute kidney injury.^{21,22}

Current research supports calcium supplementation during pregnancy to reduce the risk of pre-eclampsia onset.

Doses between 1-2 g/day are considered safe and effective and this is especially important for women with a low dietary intake of calcium, or those with predisposing factors putting them at risk of developing pre-eclampsia.^{23,24}

Kidney stone formation associated with calcium supplementation

Where once the recommendation in the management of kidney stones was dietary calcium restriction,²⁵ more recent research highlights that a dietary calcium intake greater than 500 mg/day may be protective against the formation of kidney stones by decreasing the absorption of oxalates and reducing urinary oxalate levels.² In contrast, supplemental calcium may increase the risk, which could be mitigated by taking the supplement with food, though this is currently speculative.²

Gastrointestinal effects and calcium supplementation

GI side effects may occur with high-dose calcium supplementation and are often associated with calcium carbonate.² Symptoms are mostly minor and may include an increased prevalence of abdominal pain, bloating, flatulence, severe diarrhoea and constipation.²⁶ In some instances, adverse effects from larger supplement doses can be severe and may require hospitalisation.²⁶

SUMMARY

Calcium plays a vital role in women's health beyond bones despite the high number of Australian women with inadequate dietary intake. Calcium deficiency can present in a myriad of ways with supplementation being a viable option for some..

Calcium D-glucarate supplementation provides hormonal regulation and liver support.



References

1. Beto, J.A. (2015) 'The Pole of Calcium in Human Aging', *Clinical Nutrition Research*. Korean Society of Clinical Nutrition, 4(1), pp.1-8. doi:10.7762/CNR.2015.4.11.
2. Li, K. et al. (2018) 'The good, the bad, and the ugly of calcium supplementation: a review of calcium intake on human health', *Clinical Interventions in Aging*. Dove Press, 13, pp.2443-2452. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6276611/> <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6276611/>
3. Tunçalp, Ö. et al. (2020) 'WHO recommendations on antenatal nutrition: an update on multiple micronutrient supplements', *BMJ Global Health*. BMJ Specialist Journals, 5(7), pp.1-4. Available at: <https://gh.bmj.com/content/5/7/e003375>
4. Winter, E. M. et al. (2020) 'Pregnancy and lactation, a challenge for the skeleton', *Endocrine Connections*. Bioscientifica Ltd., 9(6), pp. R143-R157. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7354730/>
5. Ji, M.-X. and Yu, Q. (2015) 'Primary osteoporosis in postmenopausal women', *Chronic Diseases and Translational Medicine*. Chinese Medical Association, 1(1), pp.9-13. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC643776/>
6. Admin. Australians falling short on calcium intake and consuming too much sodium, survey finds [Internet]. Arthritis WA, 2018 [cited 2020ct31]. Available from: <https://www.arthritiswa.org.au/news/australians-falling-short-on-calcium-intake-and-consuming-too-much-sodium-survey-finds/> <https://www.arthritiswa.org.au/news/australians-falling-short-on-calcium-intake-and-consuming-too-much-sodium-survey-finds/>
7. Roud AS, Sui Z, Rangan A, Grech A, Allman-Farinelli M. Low calcium intakes among Australian adolescents and young adults are associated with higher consumption of discretionary foods and beverages. *Nutrition*. 2018;55:56-146-53.
8. Cormick, G. and Belizán, J.M. (2019) 'Calcium Intake and Health', *Nutrients*. Multidisciplinary Digital Publishing Institute (MDPI), 11(7), pp.1-8. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6883260/> Straub, D.A. (2007) 'Calcium supplementation in clinical practice: a review of forms, doses, and indications', *Nutrition in clinical practice: official publication of the American Society for Parenteral and Enteral Nutrition*. *Nutr Clin Pract*, 22(3), pp.286-296. Available at: HYPERLINK <https://pubmed.ncbi.nlm.nih.gov/17507729/>
9. Straub, D.A. (2007) 'Calcium supplementation in clinical practice: a review of forms, doses, and indications', *Nutrition in clinical practice: official publication of the American Society for Parenteral and Enteral Nutrition*. *Nutr Clin Pract*, 22(3), pp.286-296. Available at: HYPERLINK <https://pubmed.ncbi.nlm.nih.gov/17507729/> <https://pubmed.ncbi.nlm.nih.gov/17507729/>
10. Lee, A. W. and Cho, S. S. (2015) 'Association between phosphorus intake and bone health in the NHANES population', *Nutrition Journal*. BioMed Central Ltd., 14(1), pp.1-7. Available at: <https://nutrition.biomedcentral.com/articles/10.1186/s12937-015-0017-0>
11. Trautvetter, U. et al. (2019) 'Calcium and Phosphate Metabolism, Blood Lipids and Intestinal Sterols in Human Intervention Studies Using Different Sources of Phosphate as Supplements - Pooled Results and Literature Search', *Nutrients*. Multidisciplinary Digital Publishing Institute (MDPI), 10(7), pp.1-25. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6073240/>
12. Jean Hailes for Women's Health (2020) Recommended daily calcium intake, *Nutrients*. Available at: <https://www.jeanhailes.org.au/health-a-z/healthy-living/nutrients/calcium> (Accessed: 21 June 2022).
13. Rondanelli, M. et al. (2019) 'Micronutrients Dietary Supplementation Advices for Celiac Patients on Long-Term Gluten-Free Diet with Good Compliance: A Review', *Medicina*. Multidisciplinary Digital Publishing Institute (MDPI), 55(7), pp.1-7. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6681258/>
14. Wu, H. and Pang, Q. (2017) 'The effect of vitamin D and calcium supplementation on falls in older adults: A systematic review and meta-analysis', *Der Orthopäde*. Orthopäde, 46(9), pp.729-736. Available at: <https://pubmed.ncbi.nlm.nih.gov/28718008/>
15. Gibson, M. E. S. et al. (2020) 'Where Have the Periods Gone? The Evaluation and Management of Functional Hypothalamic Amenorrhoea', *Journal of Clinical Research in Pediatric Endocrinology*. Galenos Yayinevi, 12(Suppl 1), pp.18-27. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7053439/>
16. Liu, D. et al. (2013) 'A practical guide to the monitoring and management of the complications of systemic corticosteroid therapy', *Allergy, Asthma, and Clinical Immunology: Official Journal of the Canadian Society of Allergy and Clinical Immunology*. BioMed Central, 9(1), pp.1-49. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC376515/>
17. 'Calcium D-Glucarate Monograph' (2002) *Alternative Medicine Review*, 7(4), pp.336-9. Available at: <https://altmedrev.com/wp-content/uploads/2019/02/AMR7-4-336.pdf>
18. Myung, S. K. et al. (2021) 'Calcium Supplements and Risk of Cardiovascular Disease: A Meta-Analysis of Clinical Trials', *Nutrients*. Multidisciplinary Digital Publishing Institute (MDPI), 13(2), pp.1-17. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7910980/>
19. Meyer, F., König, H. H. and Hajek, A. (2019) 'Osteoporosis, fear of falling, and restrictions in daily living: Evidence from a nationally representative sample of community-dwelling older adults', *Frontiers in Endocrinology*. Frontiers Media S.A., 10(SEP), pp.1-11. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6775197/>
20. Thanapattiwong, S. et al. (2020) 'Vitamin D supplement on prevention of fall and fracture: A Meta-analysis of Randomized Controlled Trials', *Medicine*. NLM (Medline), 99(34), pp.1-12. Available at: https://journals.lww.com/md-journal/Fulltext/2020/08210/Vitamin_D_supplement_on_prevention_of_fall_and_fracture.aspx
21. Assontsa Kafack, C. et al. (2019) 'Comparative effect of calcium supplementation on the incidence of pre-eclampsia and eclampsia among primigravid women', *Clinical Journal of Obstetrics and Gynecology*. Heighten Science Publications Corporation, 2(2), pp.145-149. Available at: <https://www.heighpubs.org/cjog/pdf/cjog-aid1038.pdf>
22. Poon, L. C. et al. (2021) 'A literature review and best practice advice for second and third trimester risk stratification, monitoring, and management of pre-eclampsia', *International Journal of Gynecology & Obstetrics*. John Wiley & Sons, Ltd, 154(S1), pp.3-31. Available at: <https://obgyn.onlinelibrary.wiley.com/doi/10.1002/ijgo.13763>
23. Kumar, A. and Kaur, S. (2017) 'Calcium: A Nutrient in Pregnancy', *Journal of Obstetrics and Gynaecology of India*. Springer, 67(5), pp.313-318. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5561751/>
24. Fox, R. et al. (2019) 'Preeclampsia: Risk Factors, Diagnosis, Management, and the Cardiovascular Impact on the Offspring', *Journal of Clinical Medicine*. Multidisciplinary Digital Publishing Institute (MDPI), 8(10), pp.1-22. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6832549/>
25. Sorensen, M. D. (2014) 'Calcium intake and urinary stone disease', *Translational Andrology and Urology*. AME Publications, 3(3), pp.1-13. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4708574/>
26. Reid, I. R., Bristow, S. M. and Bolland, M. J. (2015) 'Calcium supplements: benefits and risks', *Journal of Internal Medicine*. John Wiley & Sons, Ltd, 278(4), pp.354-368. Available at: <https://onlinelibrary.wiley.com/doi/10.1111/joim.12394>